arXiv:1207.4687v1 [astro-ph.HE] 19 Jul 2012

The extremely long period X-ray source in RCW 103: a descendant of Thorne-Żytkow Object?

X. W. Liu[1], R. X. Xu and G. J. Qiao

School of Physics and State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing 100871, China

J. L. Han

National Astronomical Observatories, Chinese Academy of Sciences, Beijing 100012, China

Z. W. Han

Yunnan Astronomical Observatory, Chinese Academy of Sciences, Kunming 650011, China

and

X. D. Li

Department of Astronomy, Nanjing University, Nanjing 210093, China

Received ______________________; accepted ______________________

---

[1] xiongwliu@163.com

ABSTRACT

The spin evolution of the compact neutron core in a Thorne-Żytkow Object (TŻO) is investigated to explore the origin of extremely long period X-ray source. It is found that the outflow would effectively take away angular momentum from the core when radiation pressure dominates the accretion process. Thus the compact core could quickly spin-down to the co-rotation period (e.g. several hours) within the massive envelope, in about $10^3 - 10^4$ years. The compact core could become an extremely long period compact star if the envelope is disrupted by some powerful bursts or exhausted via the stellar wind. The 6.67-hour periodic modulation of the central compact object (CCO) in supernova remnant RCW 103 could be naturally understood as the descendant of a TŻO.

Subject headings: ISM: supernova remnants – stars: evolution – stars: individual (1E161348-5055) – stars: neutron

1. Introduction

The central compact object (CCO) in supernova remnant RCW 103, 1E161348-5055 (hereafter 1E1613), is a very strange X-ray point source. It has a X-ray luminosity in the range of $\sim 10^{33} - 10^{35}$ erg/s, and has no identified radio, infrared or optical counterpart. In 2005, a periodic modulation of 6.67 hours was detected by the EPIC MOS cameras onboard XMM-Newton (De Luca et al. 2006), and any other periodicities with $P \geq 12$ ms are excluded with high confidence. However, RCW 103 is a supernova remnant of only about 2000 years old (Nugent et al. 1984; Carter et al. 1997)), which means that the CCO 1E1613 is very young. These features make 1E1613 a unique source among all the CCOs (De Luca 2008).

The 6.67-hour periodicity is normal if it is the orbital period of a binary. But it is difficult to explain how the companion survive from the supernova explosion and escape from the multi-wavelength observations, and why the spin period of 1E1613 has not been detected if 6.67 hour is the orbital period. If 6.67 hour is the spin period of 1E1613, it would be the longest one of all isolated pulsars. How could an isolated pulsar spin-down to such a long period in 2000 years? De Luca et al. (2006) proposed a scheme that a magnetar with an initial period of 0.3 s and a surface magnetic field of 5×10^{15} G surrounded by a disk of $3 \times 10^{-5} M_{\odot}$ may spin down to a period of 6.67 hour in 2000 years, owning to the extremely large propeller torque. This initial period is too large according to the magnetar theory (Duncan & Thompson 1992), and its current period is very different from the range of 2-12 s for the other magnetars detected as anomalous X-ray pulsars (AXPs) and soft gamma-ray repeaters (SGRs). The AXP 0142+61 (Wang et al. 2006) has a period of only 8.7 s. We noticed that even in binaries, extremely long periods have been detected from X-ray sources in binaries, such as the $\sim$ 2.7 hour period X-ray source in 2S 0114+65 (Finley, Belloni, & Cassinelli 1992; Li & van den Heuvel 1999) and the $\sim$ 1.5 hour period

X-ray source in 4U 2206+54 (Wang 2009).

On the other hand, the predicted stars with compact neutron cores remain mystery which was first proposed by Gamow (1937) and Landau (1938) to solve the stellar energy problem. The idea was abandoned when the stellar energy is realized to be generated from thermonuclear reactions (Bethe 1939). Thorne & Żytkow (1975, 1977) discussed the possible equilibrium states of a massive star with a degenerate neutron core, which was called Thorne-Żytkow Object (TŻO). A TŻO could be produced in at least three ways: 1) The expansion of the normal star in a binary drags the companion neutron star (NS) and gradually merges (Taam et al. 1978); 2) A collision of a NS with a massive main-sequence star (MS star) (Benz & Hills 1992); and 3) A supernova (SN) explosion gives the young NS an appropriate kick velocity and a direction which make it embed in its binary companion (Leonard et al. 1994). Podsiadlowski et al. (1995) estimated a total number of 20-200 TŻOs existing in the Galaxy given the characteristic lifetime of TŻO as being $10^5 - 10^6$ yr.

The TŻOs usually manifest as red supergiants or red giants (Thorne & Żytkow 1977; Eich et al 1989). But they can be distinguished from the ordinary ones since they have extraordinary high abundances of lithium and rapid proton process elements (Thorne & Żytkow 1977). The final fates of TŻOs are not entirely clear. It is generally believed that a neutron core accreted enough material and finally transform to a black hole (BH), and that the massive envelope may be dissipated and form a disk. Here we discuss another possibility. If a powerful burst occurs before the neutron core transform to a BH, the burst can disrupts the envelope, the neutron core may looks like a CCO, and the envelope may mix with the original SNR which produced the neutron core or form a new SNR. Usually, the spin energy of the neutron core is much smaller than its kinetic energy and the accretion energy, thus has little influence to the evolution of a TŻO. However, if the core could be observable after the envelope is disrupted, the spin behavior of the core could be very important. We proposed that 1E1613 and RCW 103 are the T$\dot{Z}$O descendant. We will deliberate how the compact core in a T$\dot{Z}$O spin down to a long period, and analyze different fates of the compact cores in T$\dot{Z}$Os. There may be two possible approaches for a T$\dot{Z}$O transforming to 1E1613 and RCW 103.

Case A. About 2000 years ago, the more massive star of a binary experienced a supernova explosion, which produced RCW 103 and a compact star. By some possibility the compact star got an appropriate kick velocity and direction to be embedded in the companion MS star and formed a T$\dot{Z}$O (Leonard et al. 1994). The T$\dot{Z}$O would manifest as a red giant or red supergiant afterwards (Eich et al 1989), which has angular frequency much smaller than its compact core. Thus, the core is then spun down to the co-rotational period (6.67 hour) by the envelop efficiently, as discussed below. And then a powerful burst occurred and destroyed the envelope (e.g., due to a phase transition of its crust (Cheng et al. 1998), or other processes such as thermal nuclear explosion). As a result, the envelope mixed together with the original SNR RCW 103, and the core became observable as a central compact object, i.e. 1E1613. The accretion from the residual envelope or a fall-back disk reproduces the observational X-ray emission.

Case B. More than 2000 years ago, a neutron star in a binary was embedded in its companion star or was swallowed by it (Benz & Hills 1992) and formed a T$\dot{Z}$O. The core spin down quickly to the co-rotational period, 6.67 hours. About 2000 years ago, a global thermal nuclear explosion took place or the whole neutron star occurred a phase transition as a quark-nova (Ouyed et al. 2002). The powerful energy emission destroyed the T$\dot{Z}$O envelope and formed an analogous supernova remnant, i.e. RCW 103. The central compact object 1E1613 accretes from the residual envelope or a fall-back disk and emits the observational X-ray.

In both cases above, the central compact object would has very small proper motion velocity, which is consistent with the observation of 1E1613.

2. Spin evolution of the neutron core in a TŻO

Podsiadlowski et al. (1995) estimated the spin evolution of a neutron core in a TŻO via the total angular momentum carried by the accretion material. They found a slowly spin-down process during the steady-burning phase where the accretion has an Eddington rate, and a quickly spin-up process in the neutrino runaway case where the accretion is not Eddington-limited which could spin-up the core to a typical period of ∼ 10 ms. Nevertheless, they did not consider the interaction between the neutron core magnetosphere and the TŻO envelope, and also ignored the outflow of the convection envelope. The convection passes through the whole envelope out to the photosphere (Thorne & Żytkow 1977). The outflow carries away the heat energy as well as the angular momentum, which would make the neutron core spin-down effectively. We now deliberate the spin evolution of the neutron core by considering both the interaction and the outflow.

2.1. The model

During the steady-burning phase of a TŻO, some of the envelope material is accreted to the neutron core surface. The radiation pressure opposes the gravitational force and forms an equilibrium. The difference between the inflow rate ($\dot{M}_{\rm in}$) and outflow rate ($\dot{M}_{\rm out}$) determines the rest accretion rate ($\dot{M}_{\rm acc}$), i.e.

$$\dot{M}_{\rm acc} = \dot{M}_{\rm in} - \dot{M}_{\rm out}. \tag{1}$$

In a spherical symmetry accretion case, $\dot{M}_{\rm acc}$ approximates the Eddington rate, i.e.

$$\dot{M}_{\rm acc} \approx 10^{18} R_6 {\rm g/s}, \tag{2}$$

where $R_6 = R_*/10^6$cm, where R_* is the compact star radius.

To estimate the inflow rate we consider the accretion in the steady medium, whose accretion rate is $\sim \pi R_{\rm a}^2 c_{\rm s}\rho$, where $R_{\rm a}$ is the accretion radius, $c_{\rm s}$ and ρ are the sound velocity and density at $R_{\rm a}$. The sum of the gravitational potential energy and the kinetic energy of a particle at $R_{\rm a}$ is zero. When $r < R_{\rm a}$, the kinetic energy of the particle cannot overcome the gravitation, thus the particle is captured by the core. For a neutron core surrounded by a envelope with a structure gave in Thorne & Żytkow (1977), the accretion rate in a steady medium is about 10^{21} g/s, much larger than the rest accretion rate. The inflow rate should be equal the accretion rate in a steady medium at $R_{\rm a}$, if the radiation pressure is insignificant. When considering radiation pressure, we take

$$\dot{M}_{\rm in}(R_{\rm a}) = \eta\pi {R_{\rm a}}^2 c_{\rm s}\rho, \tag{3}$$

where $0 < \eta < 1$. If we set

$$\dot{M}_{\rm in}(R_*) = \dot{M}_{\rm acc}, \tag{4}$$

we get

$$\dot{M}_{\rm out}(R_*) = 0$$

and

$$\dot{M}_{\rm out}(R_{\rm a}) = \dot{M}_{\rm in}(R_{\rm a}) - \dot{M}_{\rm acc}.$$

In the range of $R_* \leq r \leq R_{\rm a}$, the outflow can be different for different TŻOs, we take an exponential form of:

$$\dot{M}_{\rm out}(r) = (\frac{r - R_*}{R_{\rm a} - R_*})^{\alpha}(\dot{M}_{\rm in}(R_{\rm a}) - \dot{M}_{\rm acc}), \tag{5}$$

where the parameter $\alpha \sim 1$.

The above equations provide the distribution of the inflow and outflow inside $R_{\rm a}$. Their radial velocities are determined by the gravitational force, radiation pressure and the distribution. Their rotational velocities, which are related to the angular momentum, should be determined by the interaction between them and the magnetosphere of the neutron core.

The accretion in a TŻO is different from the traditional accretion of a NS. The envelope has extremely slow rotation (Podsiadlowski et al. 1995) thus a disk could not form around the neutron core. The interaction between the magnetosphere and accretion flow would not confined by the Alfvén radius (Lamb et al. 1973) since the structure of the accretion region is radiation-pressure dominant.

The magnetosphere of the neutron core is insignificant in the radial direction since the convection flows are radiation-pressure dominated. Nevertheless, it would dominate the rotational movement of the flows because the gravitational force and radiation pressure do not work in this direction. If the flows rotate so fast, that the centrifugal force can compete with the gravitational force, the structure of accretion region would be affected. If the centrifugal force is much smaller than the gravitational force, the rotational movement and radial movement should be relatively independent. Therefore, we can seperatedly study the rotational interaction between the magnetosphere and the flows while their distributions are determined by the radial direction forces.

The rotational velocities of the flows are shown in Figure 1. The flows have a potential to co-rotate with the magnetosphere with a co-rotational velocity

$$v_{\rm co} = r\Omega \tag{6}$$

because of the magnetic freezing effect, where Ω is the angular velocity of the compact core. But the magnetic field can draw the flows to a velocity at most with kinetic energy density equal to the magnetic energy density, i.e.

$$v_{\rm dr} = \beta B(4\pi\rho)^{-1/2}, \tag{7}$$

here $0 < \beta \leq 1$, and B is the magnetic field strength. Thus the real rotational velocity v_ϕ

should be the smaller one of $v_{\rm co}$ and $v_{\rm dr}$, i.e.

$$v_\phi = min(v_{\rm co}, v_{\rm dr}). \tag{8}$$

As shown in figure 1, the flows would be drawn by the magnetosphere with $v_{\rm dr}$ when they are far from the core, and they would co-rotate with the magnetosphere when they are near to the core. At a radius $R_{\rm e}$,

$$v_{\rm co}(R_{\rm e}) = v_{\rm dr}(R_{\rm e}),$$

we obtain

$$R_{\rm e} = \beta B(4\pi\rho)^{-1/2}\Omega^{-1}. \tag{9}$$

Note that B and ρ are the functions of radius r.

When a particle accesses the accretion radius $R_{\rm a}$, it will be drawn by the magnetosphere and rotate around the core with velocity $v_{\rm dr}$. If it passes through the equal velocity radius $R_{\rm e}$, it would co-rotate with the core. Therefore, the particle should be accelerated in the magnetosphere and get angular momentum from the core. If a particle finally hits to the core surface it would return the angular momentum to the core, as well as the initial angular momentum when it was in the envelope. But if it goes out it takes away angular momentum from the core, and helps the core to spin down.

If a particle goes out from $R_{\rm e} \le r < R_{\rm a}$, the angular momentum it takes away from the core is

$$j = mrv_{\rm dr}, \tag{10}$$

where m is the particle mass. If a particle goes out from $R_* < r < R_{\rm e}$, it would take away an angular momentum of

$$j = mR_{\rm e}v_{\rm co}(R_{\rm e}). \tag{11}$$

Note this value is calculated at $R_{\rm e}$ not at r, because the particle would co-rotate with the core when going out from r to $R_{\rm e}$. When the particles are going out from $R_{\rm e} \le r < R_{\rm a}$ to

$R_{\rm a}$, they could also get a bit more angular momentum from the core because the field lines are sweeping them. We ignore this additional amount of the angular momentum, because it is the same order as the initial value and can be degenerated with the model parameters (e.g. B and β).

If we assume the envelope spins with a angular velocity of $\Omega_{\rm a}$ at $R_{\rm a}$, we get the angular momentum change rate of the neutron core as

$$I\dot{\Omega} = \dot{J} = \dot{J}_{\rm in} - \dot{J}_{\rm out}, \tag{12}$$

where

$$\dot{J}_{\rm in} = \dot{M}_{\rm in}(R_{\rm a})\Omega_{\rm a}R_{\rm a}^2 \tag{13}$$

is the angular momentum change rate from inflow and $\dot{J}_{\rm out}$ the angular momentum change rate from outflow. From equations (10) and (11) we know that the angular momentums took away by the particles go out from the range of $r < R_{\rm e}$ and $r > R_{\rm e}$ are different, thus $\dot{J}_{\rm out}$ should be the sum of the contributions in the two ranges, i.e.

$$\dot{J}_{\rm out} = \dot{J}_{r<R_{\rm e}} + \dot{J}_{r>R_{\rm e}}, \tag{14}$$

where

$$\dot{J}_{r<R_{\rm e}} = \int_{R_*}^{R_{\rm e}} \nabla\dot{M}_{\rm out}v_{\rm co}(R_{\rm e})R_{\rm e}dr = \int_{R_*}^{R_{\rm e}} \frac{\alpha R_{\rm e}^2\dot{M}_{\rm out}(R_{\rm a})}{(R_{\rm a}-R_*)^{\alpha}}(r-R_*)^{\alpha-1}\Omega dr, \tag{15}$$

and

$$\dot{J}_{r>R_{\rm e}} = \int_{R_{\rm e}}^{R_{\rm a}} \nabla\dot{M}_{\rm out}v_{\rm dr}rdr = \int_{R_{\rm e}}^{R_{\rm a}} \frac{\alpha\dot{M}_{\rm out}(R_{\rm a})}{(R_{\rm a}-R_*)^{\alpha}}(r-R_*)^{\alpha-1}\beta B(4\pi\rho)^{-1/2}rdr. \tag{16}$$

In the above equations, B is defined by the structure of the magnetosphere, ρ defined by the structure of the envelope. Therefore, the spin evolution of the compact core in a TŻO could be numerically obtained from equations (2), (3), (5), (9), and (12)-(16), with a set of given parameters.

2.2. The results

In the flowing calculations, we took a dipole magnetic field for the compact core, where the magnetic moment $\mu = Br^3$. We adopted the envelope structure of TŻO given by Thorne & Żytkow (1977), with $R_{\rm a} \approx r_{\rm o-m} = 10^9$ cm, $R_* = 10^6$ cm, and $\rho = 10^{12} r^{-2}$ g/cm^3 when supposed the exponential density distribution. We took the envelope at R_a spin period $P_{\rm a} = 2\pi/\Omega_{\rm a} = 6.67$ hours (the modulation period of 1E1613) although the results are not sensitive to $P_{\rm a}$.

If the core spins too fast (e.g. Ω_1 in Fig. 1) the centrifugal force may exceed the gravitational force at a radius where the model should be invalid. When the core spins so fast that the light speed radius ($r_{\rm L} = c/\Omega$, where c is the speed of light) is smaller than the accretion radius (i.e. $r_{\rm L} < R_{\rm a}$), the upper limit of integral in equation (16) should change from $R_{\rm a}$ to $r_{\rm L}$, since the interaction between particles and magnetosphere is negligible when $r > r_{\rm L}$. On the other hand, when the core spins too slowly (e.g. Ω_3 in Fig. 1) where $R_{\rm e} > R_{\rm a}$, equation (16) does not work and the upper limit of integral in equation (15) should change from $R_{\rm e}$ to $R_{\rm a}$.

Figure 2 shows the spin evolution of a compact core in a TŻO with different core parameters and same model parameters, where P_0 is the initial period of the core, and $\mu_{30} = \mu/10^{30}$ G cm^3. It shows that a compact core can spin down to the envelope spin period $P_{\rm a}$ in about $10^3 - 10^4$ years, then co-rotate with the envelope. A core with stronger magnetic field could spin down faster than a weaker one. The spin-down timescale is approximately in proportion to the initial period.

Figure 3 shows the spin evolution of a compact core in a TŻO with same core parameters and different model parameters, where $P_0 = 0.2$ and $\mu_{30} = 1$. It shows that a core with smaller α, larger β and η spin down more effectively than the one with a larger α, smaller β and η. The spin-down timescale is approximately in inverse proportion to the accretion fraction η.

When the core has spun down it co-rotate with the T$\dot{Z}$O envelope during the steady-burning phase. If the neutrino runaway phase occurs, it could be spun up by the huge rest accretion flow. If not the neutrino runaway phase does not occur the core co-rotate until something else happens.

3. The fate of the compact core in a T$\dot{Z}$O

The compact cores in the T$\dot{Z}$Os with different initial conditions could have different fates. The relation between the initial condition and final fate is not clear. The core could transform to a BH if it accretes enough material or becomes a long period isolated compact star when the envelope vanished.

If the core evolved to a BH, its spin information should transform to the specific angular momentum of the BH, which may be tested from the X-ray properties when accreting from an disk around (Zhang et al. 1997). Usually, the BH would spin fast since it has accreted enough material with angular momentum.

It could evolve into an isolated compact star when the envelope vanishes in a relaxed or an abrupt way. A T$\dot{Z}$O usually has strong stellar wind, thereby it may lose a large part of the envelope material. The core might be naked when most of the envelope has been lost, and the core could finally evolve into an isolated neutron star with an extremely long period, which might be also detected as an X-ray source due to remnants thermal energy or accretion from the residual envelope.

Three possible processes might destroy an envelope in an abrupt way. 1) When non-steady burning occurs, it would disturb the stable evolution. As its energy is not large enough it may not disrupt the envelope, but a series of nuclear burst could speed up the progress. 2) If the strange quark matter is the real base-state of hadron (Witten 1984), the phase transition might happen in extreme conditions. The hadron material transfers to quark material, and the process releases a large amount of energy which is enough to disrupt the $\mathrm{T\dot{Z}O}$ envelope. Ouyed et al. (2002) proposed that a neutron star could transfer to a quark star, which releases an extremely large amount of energy, and appears as a quark-nova. If this happens after the neutron core has spun down, the envelope can be disrupted. The supernova explosion leaves the quark star as a central compact star with long period. 3) When the core has accreted enough material, the hadron crust might break and transform to quark material (Cheng et al. 1998). If the transforming crust is larger than $\sim 10^{-5}M_{\odot}$, the energy could be enough to disrupt the envelope, which may also result in a long period isolated pulsar surrounded by diffuse medium rather than supernova remnant. If the former supernova remnant has not disappeared, the disrupted envelope would mix with it.

The compact degenerated star, including BH, should has a small proper motion velocity since it was generated inside a $\mathrm{T\dot{Z}O}$.

4. Discussion and conclusions

The mysterious $\mathrm{T\dot{Z}O}$s should exist in our Galaxy since a neutron star has various possibility to enter into a MS star. They could be verified by direct observations of a $\mathrm{T\dot{Z}O}$ or indirect observation of a descendant. Unfortunately, no direct observation is available. To indirectly verify the existence of the $\mathrm{T\dot{Z}O}$s we need to understand what their descendants could be. In this paper, we studied the spin evolution of the compact core inside a $\mathrm{T\dot{Z}O}$, and concluded that the core can quickly spin down to the co-rotation period in the inner part of the envelope. A $\mathrm{T\dot{Z}O}$ could evolve into a long period isolated X-ray compact star, surrounded by a supernova or quark-nova remnant. The observations of 1E1613 in RCW

103 are in accord with this scheme in the long periodicity, small proper motion, and other observed X-ray emission features.

We have calculated the spin evolution of the compact core in the possible equilibrium states of a massive star with a degenerate neutron core (i.e. the TZOs). The compact core of a TŻO would spin-down quickly as braked by the interaction between the envelope and magnetosphere. The core could manifest as a long period isolated compact star if an energetic burst destroys the envelope or it is faded away with the stellar wind. The CCO in RCW 103, i.e. 1E1613, might be a descendant of a disrupted TŻO. A few compact stars possess an extremely long period. Most of them may not be observable because they do not have enough accretion from surroundings to produce strong X-ray emission.

The compact stars could also be one of the possible origin of the extremely long period objects in binary X-ray sources, if they were captured by other stars. In this scenario the extremely strong magnetic field magnetars and their magnetic field decay are not necessary, as discussed by Li & van den Heuvel (1999) for the origin of the $\sim$ 2.7 hour period X-ray source in 2S 0114+65 and by Wang (2009) for the $\sim$ 1.5 hour period X-ray source in 4U 2206+54.

We would like to thank the pulsar group of PKU for useful discussions. This work is supported by the National Natural Science Foundation of China (Grant Nos. 10935001, 10973002, 10833003), the National Basic Research Program of China (Grant Nos. 2009CB824800, 2012CB821800), and the John Templeton Foundation.

REFERENCES

Benz, W., & Hills, J. G. 1992, ApJ, 389, 546

Bethe, H. A. 1939, Phys. Rev., 55, 434

Carter, L. M., Dickel, J. R., & Bomans, D. J. 1997, PASP, 109, 990

Cheng, K. S., Dai, Z. G., Wei, D. M., & Lu, T. 1998, Science, 280, 407

De Luca, A., Caraveo, P. A., Mereghetti, S., Tiengo, A., & Bignami, G. F. 2006, Science, 313, 814

De Luca, A. 2008, AIPC, 983, 311

Duncan, R. C., & Thompson, C. 1992, ApJ, 392, L9

Eich, C., Zimmermann, M. E., & Thorne K. S. 1989, ApJ, 346, 277

Finley, J., Belloni, T., & Cassinelli, I. P. 1992, A&A, 262, L25

Gamow, G. 1937, Structure of Atomic Nuclei and Nuclear Transformations. Oxford univ. Press, Oxford

Lamb, F. K., Pethick, C. J., & Pines, D. 1973, ApJ, 184, 271

Landau, L. D. 1938, Nature, 141, 333

Leonard, P. J. T., Hills, J.G., & Dewey, R. J. 1994, ApJ, 423, L19

Li, X. D., & van den Heuvel, E. P. J. 1999, ApJ, 513, L45

Nugent, J. J., Pravdo, S. H., Garmire, G. P., Becker, R. H., Touhy, I. R., & Winkler, P. F. 1984, ApJ, 284, 612

Ouyed, R., Dey, J., & Dey, M. 2002, A&A, 390, L39

Podsiadlowski, P., Cannon, R. C., & Rees, M. J. 1995, MNRAS, 274, 485

Taam, R. E., Bodenheimer, P., & Ostriker, J. P. 1978, ApJ, 222, 269

Thorne, K. S., & Żytkow, A. N. 1975, ApJ, 199, L19

Thorne, K. S., & Żytkow, A. N. 1977, ApJ, 212, 832

Wang, W. 2009, MNRAS, 398, 1428

Wang, Z. X., Chakrabarty, D., & Kaplan, D. L. 2006, Nature, 440, 772

Witten, E. 1984, Phy. Rev. D. 30, 272

Zhang, S. N., Cui, W., & Chen, W. 1997, ApJ, 482, L155

This manuscript was prepared with the AAS LaTeX macros v5.2.



Fig. 1.— The rotational velocity of a particle in accretion region. R_* is the compact neutron core radius, $R_{\rm e}$ the equal velocity radius, and $R_{\rm a}$ the accretion radius where accretion begins. When a particle accesses the accretion radius $R_{\rm a}$, it is drawn by the magnetosphere and rotates around the core with velocity $v_{\rm dr}$. If it could passes through the radius $R_{\rm e}$, it will co-rotate with the core with velocity $v_{\rm co}$. When a particle goes out from the range of $R_{\rm e} \leq r < R_{\rm a}$ or $R_* < r < R_{\rm e}$, it could not be decelerated, thus would take away angular momentum with an amount given in equations (10) and (11), respectively.



Fig. 2.— The spin evolution of the compact core in a T$\dot{Z}$O with different core parameters and same model parameters, where $\alpha = 1$, $\beta = 1$ and $\eta = 0.5$. We take I= 10^{30}gcm^2 for the compact core and $P_{\rm E} = 6.67$ hours for the envelope.



Fig. 3.— The spin evolution of the compact core in a TŻO with same core parameters and different model parameters, where $\mu_{30} = 1$ and $P_0 = 0.2$ s. We take $I = 10^{30}\mathrm{gcm}^2$ for the compact core and $P_{\mathrm{E}} = 6.67$ hours for the envelope.